

ZELLEROMYCES GIENNENSIS SP. NOV. (RUSSULALES), A GASTEROID FUNGUS FROM THE SOUTH OF SPAIN

B. MORENO-ARROYO¹, J. GOMEZ² & F.D. CALONGE³

1. Depto. de Biología y Ecología, Facultad de Ciencias,
Universidad de Córdoba,
Avda. San Alberto Magno, s/n,
14004 Córdoba, Spain.
2. Asociación Micológica de las Sierras Subbéticas,
Mesones 4, 14800 Priego, Córdoba, Spain.
3. Real Jardín Botánico, CSIC, Plaza de Murillo, 2,
28014 Madrid, Spain.

SUMMARY : *Zelleromyces giennensis* is proposed and described ■ a new species, including data on its taxonomy, ecology, morphology, anatomy and relationships with related species.

KEY WORDS : Russulales, Elasmomycetaceae, *Zelleromyces giennensis*, taxonomy, ecology, Spain.

RESUMEN : Se propone *Zelleromyces giennensis* como especie nueva para la ciencia, y se aportan datos sobre su taxonomía, ecología, morfología, anatomía, así como semejanzas y diferencias con especies próximas.

PALABRAS CLAVE : Russulales, Elasmomycetaceae, *Zelleromyces giennensis*, taxonomía, ecología, España.

RÉSUMÉ : *Zelleromyces giennensis* est proposé comme nouvelle espèce, avec l'apport de données sur la taxonomie, l'écologie, la morphologie, l'anatomie et les relations avec les taxa proches.

MOTS-CLEFS : Russulales, Elasmomycetaceae, *Zelleromyces giennensis*, taxonomie, écologie, Espagne.

INTRODUCTION

In previous papers (Calonge & Pegler, 1998; Moreno-Arroyo *et al.* 1998a, b) has been discussed the possible delimitation of the genus *Zelleromyces* Singer & A. H. Sm., in relation to *Mariellia* Mattir. and *Gymnomyces* Massee & Rodway, and the key proposed by Beaton *et al.* (1984) is considered as the most suitable one to separate the genera of Elasmomycetaceae. The number of *Zelleromyces* species already described reaches the figure of 15, from which only two have been previously recorded in Spain, *Z. hispanicus* Calonge & Pegler (Calonge & Pegler, 1998) and *Z. meridionalis* Calonge, Moreno-Arroyo

& Gómez (Moreno-Arroyo *et al.* 1998a). The material studied here is preserved in Madrid (MA-Fungi) and in the personal senior author's herbarium (BM).

DESCRIPTION

Zelleromyces giennensis, Moreno-Arroyo, Gómez & Calonge sp. nov. (Figs. 1-4)
Etym.: *giennensis*, belonging to the province of Jaen, S. Spain.

Gasterocarpus 1-2.5 cm *latus*, *subglobosus*, *vel lobatus*, *sessilis*. *Peridium* *album*, *deinde* *alutaceis*, *laeve*, *glabrum*. *Gleba* *alba*, *deinde* *alutacea*, *loculis* *irregulariter elongatis*. *Columella* *nulla*. *Sporae* *statismosporicae*, 10-13 × 7-9 µm, *ovoideae* *vel* *ellipsoideae*, *reticulatae* *amyloideae*. *Basidia* 30-45 × 6-10 µm, *cylindrica*, 1-sporigera. *Cystidium* *nulla*. *Trama* *hymenophoralis* *cum* *hyphae* *afibulatae*, *hyalinae*. *Peridium* *cum* *elementis* *laticiferis* *instructa*. *Sphaerocystis* *carens*. *Epicutis* *in* *trichodermium*. *Peridiopellis* *pseudoprosenchymatica*. *Hispania*, *Giennen*, *La Aliseda*, *subhypogaeus*, *subter* *Pinus halepensis*, 28-II-1994, legit J. Gómez et B. Moreno, MA-Fungi 38674 (*Holotypus*); BM 213.

Gasterocarp 1-2.5 cm diam., subglobose, somewhat lobed, sessile (Fig. 1). **Peridium** whitish drying pale yellowish, smooth, matt. **Gleba** white drying yellowish, labyrinthoid, with locules irregularly arranged, empty or partially filled. **Columella** absent. **Basidiospores** 10-13 × 7-9 µm (incl. orn.), ellipsoid, rarely ovoid (Figs. 2-4), hyaline, with a myxosporium of continuous or interrupted ridges forming an incomplete reticulum (Figs. 3-4), strongly amyloid. **Basidia** 30-45 × 6-10 µm, cylindrical, 1-spored. **Sterigma** 3-5 µm long. **Subhymenium** pseudoparenchymatous. **Cystidia** absent. **Hymenophoral trama** made of hyaline hyphae, 3-6 µm diam., gelatinized, thin-walled, septate, clampless. **Peridiopellis** 150-200 µm, pseudoprosenchymatous, with gelatinized, agglutinated hyphae. **Epicutis** a trichodermium, with hyaline hyphae, 4-5 µm diam., septate, gelatinized. **Laticiferous elements**, 5-8 µm diam., only present in the peridium. **Sphaerocysts** absent.

Material studied: Spain, Jaen, La Aliseda, 600 m, subhypogaeous, under *Pinus halepensis*, 30 basidiomata forming colonies, 28-II-1994, legit J. Gómez & B. Moreno, MA-Fungi 38674 (*Holotypus*); BM 213.

DISCUSSION

Zelleromyces giennensis is a well-defined species with a series of characters not found in any other described taxon of this genus. Basidiospores subreticulate, ellipsoid; peridium with laticiferous hyphae; epicutis ■ trichodermium; basidia 1-spored, and columella, cystidia and sphaerocysts absent.

Other species with ellipsoid spores are: *Z. cinnabarinus* Singer & A. H. Sm., which has ■ cinnabar red peridium when fresh and broader spores (14-17 × 11-13 µm); *Z. oregonensis* Singer & A. H. Sm., with spiny spores (Singer & Smith, 1960). *Zelleromyces gardneri* (Zeller & Dodge) Singer & A. H. Sm., has spores similar to *Z. giennensis*, ellipsoid to ovoid, but shows a dendroid columella (Singer & Smith, 1960). Another Mediterranean species close to our material is *Z. josserandi* Malençon (Malençon, 1975), but with spores ovoid, basidia 4-spored and basidioma broader, 2-4 cm diam. The remaining species of *Zelleromyces* are remote from *Z. giennensis*, considering the taxonomic features.

Again, we have had difficulties in finding the appropriate genus for our collections, as commented upon in previous occasions (Moreno-Arroyo *et al.*, 1998a, b). However, the presence of reticulate spores, laticiferous hyphae in the peridium and absence of sphaerocysts induced us to include it within *Zelleromyces*, following Beaton *et al.* (1984) and Zhang & Yu (1990).

Zelleromyces giennensis seems to be a Mediterranean species associated with *Pinus halepensis*, being the third of a series of species recently found in Spain: *Z. hispanicus* Calonge & Pegler (Calonge & Pegler, 1998) growing under *Pinus sylvestris* and *Z. meridionalis* Calonge, Moreno-Arroyo & Gómez (Moreno-Arroyo *et al.*, 1998a) which grows under *Quercus ilex* subsp. *ballota*. It is expected that these fungi are widespread in our region and that new gasteroid Russulales will be found in future, as the attention and effort put in this subject is increased.

ACKNOWLEDGEMENTS: We express our gratitude to Dr. G. López González for critical comments and to Mr. M. Jerez for valuable SEM assistance. One of us (FDC) acknowledges the financial support received from the DGES and CSIC, under research project no. PB 95-0129-003-01.

REFERENCES

- BEATON G., PEGLER D. N. & YOUNG T. W. K., 1984 — Gasteroid Basidiomycota of Victoria State, Australia 2. Russulales. *Kew bulletin* 39: 669-698.
- CALONGE F. D. & PEGLER D. N., 1998 — *Zelleromyces hispanicus* sp. nov. (Russulales, Elasmomycetaceae), an orange-red species possibly related to *Lactarius aurantiacus*. *Cryptogamie, mycologie* 19: in press.
- MALENÇON G., 1975 — Champignons hypogènes du Nord de l'Afrique II. Basidiomycetes. *Revue de mycologie* 39: 279-306.
- MORENO-ARROYO B., GÓMEZ J. & CALONGE F. D., 1998a — *Zelleromyces meridionalis* (Russulales, Elasmomycetaceae), a new species from Spain. *Mycotaxon* (in preparation).
- MORENO-ARROYO B., GÓMEZ J. & CALONGE F. D., 1998b — *Gymnomyces dominguezii* sp. nov. (Russulales, Elasmomycetaceae), from Spain. *Mycological research* (in press).
- SINGER R. & SMITH A. H., 1960 — Studies on seotiaceous fungi. IX. The astrogastraceous series. *Mem. Torrey Bot. Club* 21: 1-112.
- ZHANG B. C. & YU Y. N., 1990 — Two new species of gasteroid Russulales from China, with notes on taxonomy of *Gymnomyces*, *Martellia* and *Zelleromyces*. *Mycological research* 94: 457-462.






Fig. 1. — *Zelleromyces giennensis*. Basidiomata showing peridium and gleba. MA-Fungi 38674.
Fig. 2. — *Zelleromyces giennensis*. Basidiospores observed under the LM, after treatment with KOH. A typical subreticulate, amyloid myxosporium can be distinguished. MA-Fungi 38674.



Figs. 3-4. — *Zelleromyces giennensis*. Basidiospores observed under SEM at different magnification to see in more detail the subreticulate myxosporium. MA-Fungi 38674.